

CORRECTION

Open Access



Correction to: High-fat diet or low-protein diet changes peritoneal macrophages function in mice

Ed Wilson Santos^{1*}, Dalila Cunha de Oliveira¹, Araceli Hastreiter¹, Jackeline Soares de Oliveira Beltran¹, Marcelo Macedo Rogero², Ricardo Ambrósio Fock¹ and Primavera Borelli¹

Correction

The original version of this article [1], published on 28 June 2016, contains a mistake. The part labels in Fig. 1 are missing. The corrected version of Fig. 1 is given below.

Author details

¹Experimental Hematology Laboratory, Department of Clinical e Toxicological Analyses, Faculty of Pharmaceutical Sciences, University of Sao Paulo, Avenida Lineu Prestes, 580 – Bloco 17, São Paulo, SP 05508-900, Brazil.

²Department of Nutrition, School of Public Health, University of Sao Paulo, Sao Paulo, Brazil.

Received: 16 February 2018 Accepted: 16 February 2018

Published online: 26 February 2018

Reference

1. Santos, et al. High-fat diet or low-protein diet changes peritoneal macrophages function in mice. 2016;41:6. <https://doi.org/10.1186/s41110-016-0006-x>.

* Correspondence: ed.wilson@usp.br

¹Experimental Hematology Laboratory, Department of Clinical e Toxicological Analyses, Faculty of Pharmaceutical Sciences, University of Sao Paulo, Avenida Lineu Prestes, 580 – Bloco 17, São Paulo, SP 05508-900, Brazil
Full list of author information is available at the end of the article






Fig. 1 Representative results of flow cytometry to cellularity of the peritoneal animal of group (a) control, (b) malnourished, and (c) high-fat; double labeled (CD11b + and F4/80+) group (d) control, (e) malnourished and (f) high-fat